


















version 2 October 2020 Nitrogen-containing compounds formula, CAS #, purity, amount, type of packaging, price in US \$	Structure	$\delta^2\text{H}$ (mean value in ‰ vs. VSMOW, $\pm 1\sigma$) (range) (# of measurements)	$\delta^{13}\text{C}$ (mean value in ‰ vs. VPDB, $\pm 1\sigma$) (range) (# of measurements)	$\delta^{15}\text{N}$ (mean value in ‰ vs. AIR, $\pm 1\sigma$) (range) (# of measurements)	$\delta^{18}\text{O}$ and $\delta^{34}\text{S}$ (mean values in ‰ vs. VSMOW or VCDT, $\pm 1\sigma$) (range) (# of measurements)	for EA for GC liquid volatile
Acetanilide #1 , $\text{C}_8\text{H}_9\text{NO}$, CAS # 103-84-4, in glass vial, 5 g US \$250, 2 g US \$150		not determined (contains exchangeable hydrogen)	-29.53 \pm 0.01 ‰ from -29.51 to -29.54 ‰ n = 6	+1.18 \pm 0.02 ‰ from +1.16 to +1.21 ‰ n = 4	not determined	
Acetanilide #2 , $\text{C}_8\text{H}_9\text{NO}$, CAS # 103-84-4, in glass vial, 2 g US \$250		not determined (contains exchangeable hydrogen)	-29.50 \pm 0.02 ‰ from -29.48 to -29.53 ‰ n = 4	+19.56 \pm 0.03 ‰ from +19.53 to +19.60 ‰ n = 7	not determined	
Acetanilide #3 , $\text{C}_8\text{H}_9\text{NO}$, CAS # 103-84-4, in glass vial, 2 g US \$250		not determined (contains exchangeable hydrogen)	-29.50 \pm 0.02 ‰ from -29.49 to -29.52 ‰ n = 4	+40.57 \pm 0.06 ‰ from +40.52 to +40.66 ‰ n = 6	not determined	
L-Alanine , $\text{C}_3\text{H}_7\text{NO}_2$, CAS # 56-41-7, produced by SI Science in Japan, 100 mg in crimp-sealed glass vial, US \$250		not determined (contains exchangeable hydrogen)	-17.93 \pm 0.02 ‰ from -17.90 to -17.96 ‰ n = 5	+43.25 \pm 0.07 ‰ from +43.16 to +43.34 ‰ n = 4	not determined	
Caffeine #1 , USGS61, $\text{C}_8\text{H}_{10}\text{N}_4\text{O}_2$, CAS # 58-08-2, $\geq 99\%$, anhydrous, 500 mg in glass vial, US \$250		+96.9 \pm 0.9 ‰ n = 53 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	-35.05 \pm 0.04 ‰ n = 114 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	-2.87 \pm 0.04 ‰ n = 93 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	not determined	
Caffeine #2 , USGS62, $\text{C}_8\text{H}_{10}\text{N}_4\text{O}_2$, CAS # 58-08-2, $\geq 99\%$, anhydrous, 500 mg in glass vial, US \$250		-156.1 \pm 2.1 ‰ n = 64 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	-14.79 \pm 0.04 ‰ n = 105 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	+20.17 \pm 0.06 ‰ n = 96 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	not determined	
Caffeine #3 , USGS63, $\text{C}_8\text{H}_{10}\text{N}_4\text{O}_2$, CAS # 58-08-2, $\geq 99\%$, anhydrous, 500 mg in glass vial, US \$250		+174.5 \pm 0.9 ‰ n = 55 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	-1.17 \pm 0.04 ‰ n = 103 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	+37.83 \pm 0.06 ‰ n = 99 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	not determined	
Collagen powder from wild-caught marine fish , USGS88, 0.5 g in glass vial, US \$250	special procedures need to be followed when using this reference material for H, O, and S isotope ratios	(+20.1 \pm 6.3 ‰ for non-exchangeable H when following USGS procedure) n = 12 (https://doi.org/10.1021/acs.jafc.0c02610)	-16.06 \pm 0.07 ‰ n = 54 (https://doi.org/10.1021/acs.jafc.0c02610)	+14.96 \pm 0.14 ‰ n = 50 (https://doi.org/10.1021/acs.jafc.0c02610)	(+15.91 \pm 0.44 ‰, +17.10 \pm 0.44 ‰ when following USGS pre-drying procedure) n = 18 n = 12 (https://doi.org/10.1021/acs.jafc.0c02610)	
Collagen powder from porcine origin , USGS89, 0.5 g in glass vial, US \$250	special procedures need to be followed when using this reference material for H, O, and S isotope ratios	(-43.7 \pm 7.8 ‰ for non-exchangeable H when following USGS procedure) n = 12 (https://doi.org/10.1021/acs.jafc.0c02610)	-18.13 \pm 0.11 ‰ n = 64 (https://doi.org/10.1021/acs.jafc.0c02610)	+6.25 \pm 0.12 ‰ n = 48 (https://doi.org/10.1021/acs.jafc.0c02610)	(+8.37 \pm 0.40 ‰, +3.86 \pm 0.56 ‰ when following USGS drying procedure) n = 20 n = 12 (https://doi.org/10.1021/acs.jafc.0c02610)	
N,N-Dimethylaniline , $\text{C}_9\text{H}_{11}\text{N}$, CAS # 121-69-7, 99%, 1.0 mL sealed under argon in glass ampoule, US \$250		-48.2 \pm 2.2 ‰ from -45.2 to -51.0 ‰ n = 5	-23.79 \pm 0.01 ‰ from -23.78 to -23.80 ‰ n = 4	-1.15 \pm 0.03 ‰ from -1.10 to -1.18 ‰ n = 4	not determined	
EDTA #2, ethylene diamine tetraacetic acid , $\text{C}_{10}\text{H}_{16}\text{N}_2\text{O}_8$, CAS # 60-00-4, 99%, 2 g in glass vial, US \$250		not determined (contains exchangeable hydrogen)	-40.38 \pm 0.01 ‰ from -40.37 to -40.38 ‰ n = 4	-0.83 \pm 0.04 ‰ from -0.78 to -0.88 ‰ n = 6	not determined	
9-Ethylcarbazole , $\text{C}_{12}\text{H}_{13}\text{N}$, $\geq 99.5\%$, CAS # 86-28-2, ≥ 200 mg in crimp-sealed glass vial, US \$250		-102.0 \pm 1.1 ‰ from -100.6 to -103.6 ‰ n = 7	-25.36 \pm 0.02 ‰ from -25.35 to -25.39 ‰ n = 5	+3.93 \pm 0.06 ‰ from +3.87 to +4.00 ‰ n = 5	not determined	
Flour from Italian millet , USGS90, 0.5 g in glass vial, US \$250	special procedures need to be followed when using this reference material for H, O, and S isotope ratios	(-13.9 \pm 2.4 ‰ for non-exchangeable H when following USGS procedure) n = 12 (https://doi.org/10.1021/acs.jafc.0c02610)	-13.75 \pm 0.06 ‰ n = 51 (https://doi.org/10.1021/acs.jafc.0c02610)	+8.84 \pm 0.17 ‰ n = 42 (https://doi.org/10.1021/acs.jafc.0c02610)	(+35.90 \pm 0.29 ‰, -15.14 \pm 0.67 ‰ when following USGS pre-drying procedure) n = 14 n = 12 (https://doi.org/10.1021/acs.jafc.0c02610)	
Flour from Vietnamese rice , USGS91, 0.5 g in glass vial, US \$250	special procedures need to be followed when using this reference material for H, O, and S isotope ratios	(-45.7 \pm 7.4 ‰ for non-exchangeable H when following USGS procedure) n = 12 (https://doi.org/10.1021/acs.jafc.0c02610)	-28.28 \pm 0.08 ‰ n = 63 (https://doi.org/10.1021/acs.jafc.0c02610)	+1.78 \pm 0.12 ‰ n = 70 (https://doi.org/10.1021/acs.jafc.0c02610)	(+21.13 \pm 0.44 ‰, -20.85 \pm 0.72 ‰ when following USGS pre-drying procedure) n = 14 n = 12 (https://doi.org/10.1021/acs.jafc.0c02610)	
L-Glutamic acid , $\geq 99.5\%$, CAS # 56-86-0, 2 g in glass vial, US \$250		not determined (contains exchangeable hydrogen)	-28.60 \pm 0.01 ‰ from -28.58 to -28.61 ‰ n = 5	-2.38 \pm 0.04 ‰ from -2.32 to -2.42 ‰ n = 4	not determined	
Glycine #1 , USGS64, $\text{C}_2\text{H}_5\text{NO}_2$, $\geq 99.5\%$, CAS # 56-40-6, 500 mg in glass vial, US \$250		not determined (contains exchangeable hydrogen)	-40.81 \pm 0.04 ‰ n = 89 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	+1.76 \pm 0.06 ‰ n = 98 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	not determined	
Glycine #2 , USGS65, $\text{C}_2\text{H}_5\text{NO}_2$, $\geq 99.5\%$, CAS # 56-40-6, 500 mg in glass vial, US \$250		not determined (contains exchangeable hydrogen)	-20.29 \pm 0.04 ‰ n = 86 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	+20.68 \pm 0.06 ‰ n = 92 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	not determined	
Glycine #3 , USGS66, $\text{C}_2\text{H}_5\text{NO}_2$, $\geq 99.5\%$, CAS # 56-40-6, 500 mg in glass vial, US \$250		not determined (contains exchangeable hydrogen)	-0.67 \pm 0.04 ‰ n = 96 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	+40.83 \pm 0.06 ‰ n = 92 (<i>Anal. Chem.</i> , 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)	not determined	
Glycine #4 , $\text{C}_2\text{H}_5\text{NO}_2$, $\geq 99.5\%$, CAS # 56-40-6, produced by SI Science in Japan, 100 mg in crimp-sealed glass vial, US \$250		not determined (contains exchangeable hydrogen)	-60.02 \pm 0.02 ‰ from -60.00 to -60.06 ‰ n = 5	-26.63 \pm 0.02 ‰ from -26.61 to -26.65 ‰ n = 3	not determined	
N-Methylpiperidine , $\text{C}_7\text{H}_{13}\text{N}$, CAS # 626-67-5, 99%, 0.5 mL sealed under argon in glass ampoule, US \$250		-179.6 \pm 1.7 ‰ from -177.8 to -181.2 ‰ n = 5	-33.73 \pm 0.02 ‰ from -33.71 to -33.75 ‰ n = 4	+0.34 \pm 0.13 ‰ from 0.17 to 0.52 ‰ n = 8	not determined	

version 2 October 2020 Nitrogen-containing compounds formula, CAS #, purity, amount, type of packaging, price in US \$	Structure	$\delta^2\text{H}$ (mean value in ‰ vs. VSMOW, $\pm 1\sigma$) (range) (# of measurements)	$\delta^{13}\text{C}$ (mean value in ‰ vs. VPDB, \pm 1 σ) (range) (# of measurements)	$\delta^{15}\text{N}$ (mean value in ‰ vs. AIR, $\pm 1\sigma$) (range) (# of measurements)	$\delta^{18}\text{O}$ and $\delta^{34}\text{S}$ (mean values in ‰ vs. VSMOW or VCDT, $\pm 1\sigma$) (range) (# of measurements)	for EA	for GC	liquid	volatile
Nicotine #1 , C ₁₀ H ₁₄ N ₂ , ≥99 %, CAS # 54-11-5, 0.25 mg nicotine in 0.5 mL hexane sealed under argon in glass ampoule, US \$250		not determined	-29.98 ± 0.01 ‰ from -29.97 to -30.00 ‰ n = 5	-5.82 ± 0.05 ‰ from -5.75 to -5.88 ‰ n = 4	not determined				
Nicotine #2 , C ₁₀ H ₁₄ N ₂ , ≥99 %, CAS # 54-11-5, 0.25 mg nicotine in 0.5 mL hexane sealed under argon in glass ampoule, US \$250		not determined	+7.72 ± 0.02 ‰ from +7.68 to +7.75 ‰ n = 7	-5.94 ± 0.15 ‰ from -5.72 to -6.18 ‰ n = 7	not determined				
Nicotine #3 , C ₁₀ H ₁₄ N ₂ , ≥99 %, CAS # 54-11-5, 0.25 mg nicotine in 0.5 mL hexane sealed under argon in glass ampoule, US \$250		not determined	-30.05 ± 0.02 ‰ from -30.03 to -30.07 ‰ n = 7	+33.62 ± 0.18 ‰ from +33.40 to +33.83 ‰ n = 7	not determined				
Nicotine #4 , C ₁₀ H ₁₄ N ₂ , ≥99 %, CAS # 54-11-5, 0.25 mg nicotine in 0.5 mL hexane sealed under argon in glass ampoule, US \$250		not determined	-2.06 ± 0.02 ‰ from -2.04 to -2.08 ‰ n = 5	+15.49 ± 0.13 ‰ from +15.31 to +15.68 ‰ n = 7	not determined				
Nicotine #5 , C ₁₀ H ₁₄ N ₂ , ≥99 %, CAS # 54-11-5, 0.25 mg nicotine in 0.5 mL hexane sealed under argon in glass ampoule, US \$250		-161.3 ± 1.7 ‰ from -159.2 to -164.6 ‰ n = 10	-29.63 ± 0.01 ‰ from -29.61 to -29.65 ‰ n = 5	-6.03 ± 0.04 ‰ from -5.97 to -6.08 ‰ n = 5	not determined				
L-Phenylalanine , C ₉ H ₉ NO ₂ , ≥99.5 %, CAS # 63-91-2, produced by SI Science in Japan, 100 mg in crimp-sealed glass vial, US \$250		not determined (contains exchangeable hydrogen)	-11.20 ± 0.02 ‰ from -11.19 to -11.23 ‰ n = 6	+1.70 ± 0.06 ‰ from +1.64 to +1.77 ‰ n = 5	not determined				
L-Proline , C ₅ H ₉ NO ₂ , ≥99.5 %, CAS # 147-85-3, 100 mg in crimp-sealed glass vial, US \$250		not determined (contains exchangeable hydrogen)	-12.47 ± 0.01 ‰ from -12.45 to -12.49 ‰ n = 5	-7.84 ± 0.04 ‰ from -7.77 to -7.88 ‰ n = 5	not determined				
Pyrazine , C ₄ H ₄ N ₂ , CAS # 290-37-9, at least 20 mg in sealed glass capillary, US \$250		-31.8 ± 1.7 ‰ from -29.4 to -34.2 ‰ n = 6	not determined	+1.39 ± 0.04 ‰ from +1.34 to +1.43 ‰ n = 4	not determined				
N,N,N',N'-Tetra-n-butylurea , C ₁₇ H ₃₀ N ₂ O, CAS # 4559-86-8, 97 %, at least 10 mg sealed in glass capillary, US \$250		-112.4 ± 2.1 ‰ from -110.5 to -114.3 ‰ n = 4	-29.37 ± 0.02 ‰ from -29.35 to -29.40 ‰ n = 4	-5.06 ± 0.04 ‰ from -5.00 to -5.09 ‰ n = 4	not determined				
N,N,N',N'-Tetramethylurea , C ₄ H ₁₂ N ₂ O, CAS # 632-22-4, 99 %, 1.0 mL sealed under argon in glass ampoule, US \$250		-77.8 ± 0.7 ‰ from -76.7 to -78.4 ‰ n = 5	-36.24 ± 0.01 ‰ from -36.23 to -36.25 ‰ n = 4	-1.60 ± 0.04 ‰ from -1.55 to -1.64 ‰ n = 4	not determined				
Urea #1 , CH ₄ N ₂ O, ≥99.5 %, CAS # 57- 13-6, 2 g in glass vial, US \$250		not determined (contains exchangeable hydrogen)	-34.13 ± 0.03 ‰ from -34.17 to -34.09 ‰ n = 6	+0.26 ± 0.03 ‰ from +0.20 to +0.28 ‰ n = 7	not determined				
Urea #2a , CH ₄ N ₂ O, ≥99.5 %, CAS # 57- 13-6, 2 g in glass vial, US \$250		not determined (contains exchangeable hydrogen)	-9.14 ± 0.02 ‰ from -9.11 to -9.17 ‰ n = 10	+20.73 ± 0.04 ‰ from +20.67 to +20.78 ‰ n = 9	not determined				
Urea #3a , CH ₄ N ₂ O, ≥99.5 %, CAS # 57- 13-6, 2 g in glass vial, US \$250		not determined (contains exchangeable hydrogen)	+5.89 ± 0.03 ‰ from +5.85 to +5.93 ‰ n = 5	+42.05 ± 0.03 ‰ from +42.02 to +42.10 ‰ n = 5	not determined				
USGS88, marine collagen powder from wild-caught fish , 0.5 g in glass vial, US \$250	special procedures need to be followed when using this reference material for H, O, and S isotope ratios	+20.1 ± 6.3 ‰ for non- exchangeable H when following USGS procedure) n = 12 (https://dx.doi.org/10.1021/acs.jafc.0c02 2610)	-16.06 ± 0.07 ‰ n = 54 (https://dx.doi.org/10.1021/acs.jafc.0c02 610)	+14.96 ± 0.14 ‰ n = 50 (https://dx.doi.org/10.1021/acs.jafc.0c02 610)	(+15.91 ± 0.44 ‰ +17.10 ± 0.44 ‰ when following USGS pre-drying procedure) n = 18 n = 12 (https://dx.doi.org/10.1021/acs.jafc.0c02 2610)				
USGS89, porcine collagen powder , 0.5 g in glass vial, US \$250	special procedures need to be followed when using this reference material for H, O, and S isotope ratios	(-43.7 ± 7.8 ‰ for non- exchangeable H when following USGS procedure) n = 12 (https://dx.doi.org/10.1021/acs.jafc.0c02 2610)	-18.13 ± 0.11 ‰ n = 64 (https://dx.doi.org/10.1021/acs.jafc.0c02 610)	+6.25 ± 0.12 ‰ n = 48 (https://dx.doi.org/10.1021/acs.jafc.0c02 610)	(+8.37 ± 0.40 ‰ +3.86 ± 0.56 ‰ when following USGS pre-drying procedure) n = 20 n = 12 (https://dx.doi.org/10.1021/acs.jafc.0c02 2610)				
USGS90, millet flour from Italy , 0.5 g in glass vial, US \$250	special procedures need to be followed when using this reference material for H, O, and S isotope ratios	(-13.9 ± 2.4 ‰ for non- exchangeable H when following USGS procedure) n = 12 (https://dx.doi.org/10.1021/acs.jafc.0c02 2610)	-13.75 ± 0.06 ‰ n = 51 (https://dx.doi.org/10.1021/acs.jafc.0c02 610)	+8.84 ± 0.17 ‰ n = 42 (https://dx.doi.org/10.1021/acs.jafc.0c02 610)	(+35.90 ± 0.29 ‰ -15.14 ± 0.67 ‰ when following USGS pre-drying procedure) n = 14 n = 12 (https://dx.doi.org/10.1021/acs.jafc.0c02 2610)				
USGS91, rice flour from Vietnam , 0.5 g in glass vial, US \$250	special procedures need to be followed when using this reference material for H, O, and S isotope ratios	(-45.7 ± 7.4 ‰ for non- exchangeable H when following USGS procedure) n = 12 (https://dx.doi.org/10.1021/acs.jafc.0c02 2610)	-28.28 ± 0.08 ‰ n = 63 (https://dx.doi.org/10.1021/acs.jafc.0c02 610)	+1.78 ± 0.12 ‰ n = 70 (https://dx.doi.org/10.1021/acs.jafc.0c02 610)	(+21.43 ± 0.44 ‰ -20.63 ± 0.72 ‰ when following USGS pre-drying procedure) n = 14 n = 12 (https://dx.doi.org/10.1021/acs.jafc.0c02 2610)				
L-Valine #1 , USGS73, C ₆ H ₁₁ NO ₂ , CAS # 516-06-3, 99 %, 500 mg in glass vial, US \$250		not determined (contains exchangeable hydrogen)	-24.03 ± 0.04 ‰ n = 130 (Anal. Chem., 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem. 5b04392)	-5.21 ± 0.05 ‰ n = 91 (Anal. Chem., 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem. 5b04392)					
L-Valine #2 , USGS74, C ₆ H ₁₁ NO ₂ , CAS # 516-06-3, 99 %, 100 mg in glass vial, freeze-dried, US \$250		not determined (contains exchangeable hydrogen)	-9.30 ± 0.04 ‰ n = 94 (Anal. Chem., 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem. 5b04392)	+30.19 ± 0.07 ‰ n = 68 (Anal. Chem., 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem. 5b04392)					

version 2 October 2020 Nitrogen-containing compounds formula, CAS #, purity, amount, type of packaging, price in US \$	Structure	$\delta^2\text{H}$ (mean value in ‰ vs. VSMOW, $\pm 1\sigma$) (range) (# of measurements)	$\delta^{13}\text{C}$ (mean value in ‰ vs. VPDB, $\pm 1\sigma$) (range) (# of measurements)	$\delta^{15}\text{N}$ (mean value in ‰ vs. AIR, $\pm 1\sigma$) (range) (# of measurements)	$\delta^{18}\text{O}$ and $\delta^{34}\text{S}$ (mean values in ‰ vs. VSMOW or VCDT, $\pm 1\sigma$) (range) (# of measurements)	for EA for GC liquid volatile
L-Valine #3, USGS75 , $\text{C}_6\text{H}_{11}\text{NO}_2$, CAS # 516-06-3, 99 %, 100 mg in glass vial, freeze-dried, US \$250		not determined (contains exchangeable hydrogen)	$+0.49 \pm 0.07$ ‰ n = 23 <i>(Anal. Chem., 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)</i>	$+61.53 \pm 0.14$ ‰ n = 29 <i>(Anal. Chem., 2016, 88, 4294, http://dx.doi.org/10.1021/acs.analchem.5b04392)</i>		